

Cherry SWEET ARYANA® PA1UNIBO* SWEET LORENZ® PA2UNIBO*



SWEET ARYANA® PA1UNIBO*

Tree: high vigor, open canopy, good branching, early bearing, high and constant cropping.

Blossoming time: early bloom, 2-3 days pre B.Burlat, self –fertile (allelic profile: S3S4').

Fruits: medium-large size (average dominant size 28-30 mm); symmetrical depressed and heart-shaped, shiny tending to dark red skin (5 in CTIFL color scale), shiny dark reddish blackish skin at advanced ripening (6 in CTIFL color scale); firm flesh; sweet, good taste; medium long peduncle.

Harvesting season: early season; 3 days post B.Burlat; uniform ripening; optimal and long ontree fruit lifespan.



SWEET LORENZ® PA2UNIBO*

Tree: high vigor, open canopy, standard fruiting habitus, good branching, early bearing, high and constant cropping.

Blossoming time: intermediate bloom, 1 day after Burlat, self-sterile (allelic profile: S3S4).

Fruits: very large size (average dominant size 32-34 mm); symmetrical heart-shaped, shiny red skin (4 in CTIFL color scale) tending to shiny purplish red skin at advanced ripening (5-6 in CTIFL color scale); firm flesh; sweet, good taste; medium long peduncle.

Harvesting season: mid-early season, 6 days post B.Burlat, uniform ripening, optimal and long on-tree fruit lifespan.

Recommended pollenisers: Sweet Aryana[®] PA1UNIBO*, Marysa[®] PA6UNIBO*, Sweet Gabriel[®] PA3UNIBO*, Vera[◊].



* Registered Community design - Applicant: Alma Mater Studiorum - University of Bologna. ®Community Trademark- Owner: Alma Mater Studiorum - University of Bologna.